



octenisept®

Aqueous wound and mucous membrane antiseptic

Our Plus

- broad spectrum of antiseptic efficacy
- fast onset of action (1 min.) and long-term effect (24 h remanence)
- very good skin and mucous membrane tolerance
- suitable for infants and premature babies¹
- safe use during pregnancy
- painless and colourless treatment

Application areas

Wound treatment

Antiseptic treatment of traumatic, acute, chronic, surgical and burn wounds.

Mucous membrane antiseptic

- prior to diagnostic and surgical interventions in the anogenital (e.g. before obliteration of haemorrhoids), the urogenital area (e.g. before placing an intra-uterine device (IUD), the vaginal area, before prenatal, intranatal and postnatal manipulations) and in the oral cavity (e.g. before tooth extractions)
- before placing urinary tract catheters
- for preop. skin antiseptics in the area close to mucous membranes

MRSA

- for antimicrobial whole body wash with MRSA colonised patients^{2,3}

Instructions for use

- Moisturize the treated area evenly and thoroughly with the antiseptic.
- Swab method: Rub concerned areas with saturated swabs. Swabbing is the preferred method of application.
- Spray method: In individual cases spray octenisept® directly on poorly accessible areas of the skin and mucous membrane. Make sure all areas are evenly moistened.

- octenisept® may also be applied by rinsing the oral cavity

Microbiological efficacy

Efficacy	Concentration	Contact time
bactericidal	ready-to-use	30 sec.
Chlamydia trachomatis	ready-to-use	30 sec.
MRSA	ready-to-use	30 sec.
Neisseria gonorrhoeae	ready-to-use	30 sec.
Pseudomonas aeruginosa	ready-to-use	30 sec.
Gardnerella vaginalis	ready-to-use	30 sec.
Candida albicans	ready-to-use	2 min.
enveloped viruses (incl. HIV, HBV, HCV)	ready-to-use	30 sec.
Trichomonads spp.	ready-to-use	1 min.

Application area	Concentration	Contact time
ano- and urogenital	ready-to-use	1 min.
before bladder catheterisation	ready-to-use	1 min.
dabbing of the oral cavity	ready-to-use	1 min.
rinsing of the oral cavity	ready-to-use	20 sec.
skin antiseptics before caesarean operation	ready-to-use	2 min.
wounds	ready-to-use	1 min.



Product data

Composition:

100g solution contains: Octenidine dihydrochloride 0.1g, 2-Phenoxyethanol (Ph.Eur.) 2.0g

Chemical-physical data

Color	colourless
Density	ca. 1,005 g/cm ³ / 20 °C
Flash point	Not applicable
Form	liquid
pH	ca. 6 / 20 °C / concentrate

Special advice

- **To prevent possible tissue injury, the product must not be injected or applied to tissues with pressure. Adequate drainage from wound cavities must be provided (e.g. for flexible drain tube).**
- octenisept® should not be used for irrigating the abdominal cavity (e.g. intraoperatively), the urinary bladder and nose or eardrum.
- Do not swallow octenisept®.
- Do not enter it into the blood circuit, e.g. by being accidentally injected.
- Do not mix octenisept® with other compounds.
- Do not use octenisept® in combination with PVP-iodine based antiseptics.
- Bandages and incision foils can be applied after octenisept® has dried off completely.
- In rare cases octenisept® may cause slight burning.
- octenisept® can be heated up to body temperature.
- octenisept® expires three years after opening.
- Once the container has been opened octenisept® should not be used for more than three years but not beyond the expiry date.
- As a general principle: administrating of any pharmaceuticals within the first trimester of pregnancy should be carried out under strict indication and medical supervision.
- Microbiological efficacies on specific germs have only been carried out in in-vitro tests.

Information for order

Item	Delivery form	Item no.
octenisept colourless -CH- 50 ml FL	20/Carton	on request
octenisept 500 ml FL	20/Carton	on request
octenisept Anstaltspack. 500 ml FL	20/Carton	on request
octenisept 1 l FL	10/Carton	on request
octenisept 250 ml FL	10/Carton	on request
octenisept mit Sprühpumpe 250 ml FL	10/Carton	on request
octenisept Anstaltspack. 250 ml FL	10/Carton	on request
octenisept Anstaltspack. 1 l FL	10/Carton	on request
octenisept AP mit Sprühpumpe 250 ml FL	10/Carton	on request

These products are not available in every country. For more information please contact our local subsidiary or distributor.

Environmental information

schülke manufactures products economically and with advanced, safe and environmentally friendly production processes while at the same time maintaining out high quality standards.

Expert opinion and information

Please visit our website for an overview of all available literature/reports on the product: www.schuelke.com.

For individual questions:

Customer Care

Phone: +49 40 52100-666

E-Mail: info@schuelke.com

¹Krishna B.V.S., Gibb A.P. (8.1.2010): Use Of Octenidine Dihydrochloride In Meticillin-Resistant StaphylococcusAureus Decolonisation Regimens: A Literature Review, Journal of Hospital Infection, 74, 199-203

²Briese et al. (2010): Efficacy and tolerability of a local acting antiseptic agent in the treatment of vaginal dysbiosis during pregnancy; in Arch Gynecol Obstet

³There is no sufficient experience for the application on premature infants and neonates with immature skin (e.g. restricted barrier function of the skin).

octenisept®

Composition: 100 g solution contain: octenidine hydrochloride 0.1 g, phenoxyethanol (Ph.Eur.) 2.0 g ; Other ingredients: (3-amidopropyl cocoate) dimethylammonium acetate, sodium D gluconate, glycerol 85%, sodium chloride, sodium hydroxide, purified water.

- **Indications:** For repeated, short-term antiseptic treatment of mucous membranes, adjacent skin and as adjuvant antiseptic wound treatment. octenisept® is intended for superficial application and must not be applied e.g. by syringe into the depths of the tissue.

- **Contraindications:** octenisept® may not be used in cases of hypersensitivity to any of the components of the preparation. octenisept® should not be used for rinsing the abdominal cavity (e.g. intra-operatively) or the bladder, nor the tympanic membrane. - **Undesirable effects:** In rare cases transient signs of local intolerance such as a slight burning sensation, redness or itching may occur at the application site. Rinsing of the oral cavity may cause a transitory bitter sensation. - **Special warnings and special precautions for use:** Do not swallow octenisept® and do not allow octenisept® to pass into the circulation, e.g. as a result of accidental injection. To prevent possible tissue injury, the product must not be injected or applied to tissues with pressure. Adequate drainage from wound cavities must be provided (e.g. for flexible drain tube). If any of the side effects gets serious, or if you notice any side effects not listed in this user information, please tell your doctor or pharmacist.

To prevent possible tissue injury, the product must not be injected or applied to tissues with pressure. Adequate drainage from wound cavities must be provided (e.g. for flexible drain tube).

Manufacturer: Schülke & Mayr GmbH, 22840 Norderstedt, Germany, Tel. +49 40 52100-0, info@schuelke.com



Schülke & Mayr GmbH holds a Manufacturer's Authorisation according to sect 13 para 1 German Drug Law and Certificates of GMP Compliance for medicinal products.



A company of the Air Liquide-Group

schülke Headquarters

Schülke & Mayr GmbH
Robert-Koch-Str. 2
22851 Norderstedt, Germany
Phone +49 (0) 40 - 52100 - 0
Fax +49 (0) 40 - 52100 - 318
www.schuelke.com
mail@schuelke.com

Schülke & Mayr UK Ltd.
1 Jenkin Road
GB-Sheffield S9 1 AT
Phone +44 (0) 1142 - 5435 - 00
Fax +44 (0) 1142 - 5435 - 01
www.schulke.co.uk
mail.uk@schuelke.com

Schülke & Mayr Ges.m.b.H.
Seidengasse 9
1070 Wien, Austria
Phone +43 (0) 1 - 5232501-0
Fax +43 (0) 1 - 5232501-60
www.schuelke.at
office.austria@schuelke.com