

Kinnitas:	Katrin Reimand	Osakonnajuhataja	Info Folderitis
Koostas:	Kaja Vaagen	Vanemarst-õppejõud	30.01.2024
	Ees- ja perekonnanimi	Ametikoht	kuupäev

REFERENCE VALUES FOR CLINICAL CHEMISTRY TESTS

Analyte	Age	Reference range	Units	Reference
Adrenocorticotrophic hormone (P-ACTH)	≥ 18 y	Morning 7.00–10.00 a.m 1.6–13.9	pmol/L	1, 2
Alanine aminotransferase (S,P-ALAT)	< 1 y 1 y – < 4 y 4 y – < 7 y 7 y – < 13 y 13 y – < 18 y ≥ 18 y	< 71 < 31 < 36 < 44 < 45 M < 50 F < 35	U/L	1 1, 2
Albumin (S,P-Alb)	< 4 d 4 d – < 14 y 14 y – < 18 y ≥ 18 y	28–44 38–54 32–45 35–52	g/L	2, 5 2
Albumin (high sensitivity) (S,P-Alb-hs)	< 4 d 4 d – < 14 y 14 y – < 18 y ≥ 18 y	28–44 38–54 32–45 35–52	g/L	2
Albumin in cerebrospinal fluid CSF-Alb CSF-Alb/S-Alb-hs	≥ 18 y < 1 m 1 m – < 6 m 6 m – < 16 y 16 y – < 41 y 41 y – < 61 y	110–350 < 0.025 < 0.015 < 0.005 < 0.007 < 0.008	mg/L	2 1 (2004) 1, 2
Albumin in urine U-Alb/U-Crea	< 1 m 1 m – < 1 y 1 y – < 6 y 6 y – < 11 y 11 y – < 16 y ≥ 16 y	< 21 < 3.8 < 3.3 < 2.7 < 2.1 M < 2.5 F < 3.5	g/molL	1
dU-Alb	All age groups	< 30	mg/d µg/min	5 2
Alcohol surrogates (P-Alcohol surrogates)	All age groups	< 0,02 < 0,2 < 0,01 < 0,02 < 0,01 < 0,06 < 0,1	g/L	
Aldosterone (S,P-Aldo)	4 d – < 8 d 1 m – < 1 y 1 y – < 2 y 2 y – < 10 y 10 y – < 15 y	supine 5.0–175.0 supine 5.0–90.0 supine 7.0–54.0 upright 5.0–80.0 upright 4.0–48.0	ng/dL	5

	≥ 15 y	upright 3.7–43.2		42
Amylase (S,P-Amyl)	≥ 18 y	28–100	U/L	1, 2
Alpha-1-antitrypsin (S,P-AAT)	< 1 m 1 m – < 7 m 7 m – < 3 y 3 y – < 20 y ≥ 20 y	1.24–3.48 1.11–2.97 0.95–2.51 1.10–2.80 0.90–2.00	g/L	1 2
Alpha-fetoprotein (S-AFP)	< 1 m 1 m – < 6 m 6 m – < 1 y 1 y – < 19 y ≥ 19 y	> 1004 39.8 – > 1004 2.9–57.3 ≤ 5.8 ≤ 5.8		44 2
Alkaline phosphatase (S,P-ALP)	< 15 d 15 d – < 1 y 1 y – < 10 y 10 y – < 13 y 13 y – < 15 y 15 y – < 17 y 17 y – < 19 y ≥ 19 y	83–248 122–469 142–335 129–417 M 116–468 F 57–254 M 82–331 F 50–117 M 55–149 F 45–87 M 40–129 F 35–104	U/L	 2
Alkaline phosphatase, isoenzymes, fraction activity (S-ALP-isoE)	≥ 18y	liver 1 < 71 bone < 69 liver 2 < 13 intestine < 13	U/L	
Amphetamines in urine (U-Amp)	All age groups	negative		
Amikacin (S,P-Amic)	All age groups	Pre-dose (trough) concentration: therapeutic range 5–10 toxic > 10 Peak concentration therapeutic 20–25 toxic > 35	mg/L	2
Ammonia (P-NH4)	< 2 d 2 d – < 6 d 6 d – < 18 y ≥ 18 y	< 144 < 134 < 48 M 16–60 F 11–51	µmol/L	1 (2004) 2
Androstenedione (S,P-Androst)	< 14 d 14 d – < 1 y 1 y – < 6 y 6 y – < 10 y 10 y – < 12 y 12 y – < 15 y 15 y – < 19 y ≥ 19 y	< 2.54 0.09–2.1 0.09–0.57 0.2–0.92 < 2.54 M 0.51–2.01 F 0.74–6.01 M 0.87–3.55 F 0.48–6.45 M 0.98–5.32 F 1.71–4.58	nmol/L	44 2
Angiotensin-converting enzyme (S,P-ACE)	6 m – < 18 y ≥ 18 y	29–112 20–70	U/L	17
Anti-Müller Hormone (S,P-AMH)	≥ 18 y 20 y – < 25 y 25 y – < 30 y	M 0.8–14.5 F 1.2–11.7 F 0.9–9.9	µg/L	2

	30 y – < 35 y 35 y – < 40 y 40 y – < 45 y 45 y – < 51 y	F 0.6–8.1 F 0.1–7.5 F 0.03–5.5 F 0.01–2.7		
Antistreptolysin O (S,P-ASO)	< 6 y 6 y – < 18 y ≥ 18 y	< 150 < 240 < 200	kU/L	1 1, 2
Aripiprazole and dehydroaripiprazole (P-Aripiprazole+dehydroaripiprazole)	All age groups	Therapeutic range 150–500 Toxic > 1000	µg/L	11
Aspartate aminotransferase (S,P-ASAT)	< 2 d 2 d – < 6 d 6 d – < 1 y 1 y – < 4 y 4 y – < 7 y 7 y – < 13 y 13 y – < 18 y ≥ 18 y	< 122 < 110 < 96 < 71 < 53 < 50 < 46 M < 50 F < 35	U/L	1 1, 2
Barbiturates in urine (U-Bar)	All age groups	negative		
Benzodiazepines in urine (U-Bzd)	All age groups	negative		
Beta-2-microglobulin (S,P- β2-M)	1 d – < 1 m 1 m – < 6 m 6 m – < 1 y 1 y – < 4 y 4 y – < 7 y 7 y – < 10 y 10 y – < 13 y 13 y – < 16 y 16 y – < 19 y 19 y – 60 y ≥ 60 y	M 1603–4790 F 1722–4547 M 1423–3324 F 1024–3774 M 897–3095 F 999–2282 M 827–2228 F 742–2396 M 567–2260 F 546–2170 M 772–1712 F 736–1766 M 699–1836 F 704–1951 M 681–1954 F 787–1916 M 724–1874 F 555–1852 800–2400 ≤ 3000	µg/L	3 2
Betahydroxybutyrate (POCT) (B-BHB POCT)	All age groups	< 0,6	mmol/L	45
Bile acids (S,P-TBA)	≥ 18 y	< 10	µmol/L	2
Bilirubin (S,P-Bil)	< 2 d (full term) 2 d – < 3 d (full term) 3 d – < 4 d (full term) 4 d – < 7 d (full term)	< 150 < 193 < 217 < 216 < 140 < 205 < 410	µmol/L	1

	< 2 d (preterm) 2 d – < 3 d (preterm) 3 d – < 6 d (preterm) 1 m – < 18 y ≥ 18 y	< 17 < 21		2
Bilirubin (conjugated) (S,P-Bil-conj)	< 1 m ≥ 1 m	< 10 ≤ 3.4	µmol/L	1 2
B-type natriuretic propeptide, N-terminal fragment (S,P-NT-proBNP)	1 y – < 2 y 2 y – < 6 y 6 y – < 18 y ≥ 17 y	< 400 < 300 < 160 < 125 (cut-off value for excluding chronic heart failure) < 300 (cut-off value for excluding acute heart failure)	ng/L	25 2
Cocaine in urine (U-Coc)	All age groups	negative		
C-peptide (S,P-C-pept)	≥ 18 y	0.37–1.47	nmol/L	1, 2
C-reactive protein (S,P-CRP)	≥ 18 y	< 5	mg/L	1, 2
C-reactive protein, high sensitivity (S,P-CRP-hs)	< 3 w 2 m – < 16 y ≥ 18 y	< 4.1 < 2.8 For cardiovascular disease risk assessment: low risk < 1.0 medium risk 1.0–3.0 high risk > 3.0	mg/L	1 2
Dehydroepiandrosterone sulfate (S,P-DHEAS)	< 1 w 1 w – < 1 m 1 m – < 1 y 1 y – < 5 y 5 y – < 10 y 10 y – < 15 y 15 y – < 20 y 20 y – < 25 y 25 y – < 35 y 35 y – < 45 y 45 y – < 55 y 55 y – < 65 y 65 y – < 75 y ≥ 75 y	2.93–16.5 0.86–11.7 0.09–3.35 0.01–0.53 0.08–2.31 M 0.66–6.70 F 0.92–7.60 M 1.91–13.4 F 1.77–9.99 M 5.73–13.4 F 4.02–11.0 M 4.34–12.2 F 2.68–9.23 M 2.41–11.6 F 1.65–9.15 M 1.20–8.98 F 0.96–6.95 M 1.40–8.01 F 0.51–5.56 M 0.91–6.76 F 0.26–6.68 M 0.44–3.34 F 0.33–4.18	µmol/L	1, 2

Delta amino-levulinic acid in urine (U-DALA)	≥ 18 y	< 34,3	µmol/L	31
Delta amino-levulinic acid/creatinine in urine (U-DALA/U-Crea)	≥ 18 y	< 3,9	mmol/mol	36
Digoxin (S,P-Digox)	≥ 18 y	Therapeutic level 0.5–0.8 Toxic > 1.2	µg/L	51
Erythropoietin (S,P-EPO)	1 y – < 4 y 4 y – < 7 y 7 y – < 10 y 10 y – < 13 y 13 y – < 16 y 16 y – < 18 y ≥ 18 y	M 1.7–17.9 F 2.1–15.9 M 3.5–21.9 F 2.9–8.5 M 1.0–13.5 N 2.1–8.2 M 1.0–14.0 N 1.1–9.1 M 2.2–14.4 N 3.8–20.5 M 1.5–15.2 N 2.0–14.2 4.3–29.0	U/L	3 48
Ethanol (S,P-EtOH)	All age groups	< 0.2	g/L	23
Ecstasy in urine (U-Ecs)	All age groups	negative		
Phenobarbital (S,P-Phenobarb)	All age groups	Therapeutic range 10–30 Toxic > 40	mg/L	2
Ferritin (S,P-Fer)	< 1 y 1 y – < 4 y 4 y – < 7 y 7 y – < 13 y 13 y – < 18 y 18 y – < 61 y	12–327 6–67 4–67 M 14–124 F 7–84 M 14–152 F 13–68 M 30–400 F 13–150	µg/L	1 2
Folate (S,P-Fol)	< 7 y 7 y – < 12 y 12 y – < 18 y 18 y – < 66 y	> 17.3 > 37.9 > 17.8 8.8–60.8	nmol/L	44 2
Follicle stimulating hormone (S,P-FSH)	< 1 y 1 y – < 9 y 9 y – < 12 y 12 y – < 18 y ≥ 18 y	M 0.1–3.2 N 1.6–19 M 0.2–2.1 N 0.7–5.8 M 0.4–4.2 N 0.5–7.6 M 0.9–7.1 N 0.9–9.1 M 1.5–12.4 F follic. 3.5–12.5 ovul 4.7–21.5 luteal 1.7–7.7 postmenop 25.8–134.8	U/L	44 1, 2
Phosphate (S,P-P)	<15 d 15 d – < 1 y 1 y – < 5 y 5 y – < 13 y 13 y – < 16 y 16 y – < 19 y ≥ 19 y	1.71–3.15 1.47–2.54 1.33–2.06 1.28–1.82 F 1.00–1.70 M 1.11–1.88 0.94–1.55 0.81–1.45	mmol/L	44 2
Phosphate in urine U-P (first morning urine) dU-P U-P/U-Crea	≥ 18 y 12 y – < 61 y 6 m – < 1 y 1 y – < 2 y 2 y – < 3 y	13–44 13–42 1.2–19 1.2–14 1.2–12	mmol/L mmol/d mol/mol	1, 2 1, 2 4

	3 y – < 5 y 5 y – < 7 y 7 y – < 10 y 10 y – < 14 y 14 y – < 18 y	1.2–8.0 1.2–5.0 1.2–3.6 0.8–3.2 0.8–2.7		
Gamma glutamyltransferase (S,P-GGT)	< 2 d 2 d – < 6 d 6 d – < 7 m 7 m – < 1 y 1 y – < 4 y 4 y – < 7 y 7 y – < 13 y 13 y – < 18 y ≥ 18 y	< 151 < 185 < 204 < 34 < 18 < 23 < 17 M < 45 F < 33 M < 60 F < 40	U/L	1 1, 2
Gamma-hydroxybutyrate in urine (U-GHB)	All age groups	negative		
Gastrin (S,P-Gastr)	≥ 18 y	6.2–54.8	pmol/L	48
Gentamicin (S,P-Genta)	All age groups	Pre-dose (trough) concentration: therapeutic range 0.5–2 toxic > 2	mg/L	2
Glucose in serum/plasma, fasting (fS,fP-Gluc)	< 2 d 2 d – < 1 m 1 m – < 18 y ≥ 18 y	2.2–3.3 2.8–4.4 3.3–5.6 ≤ 6.0	mmol/L	5 15
Glucose in cerebrospinal fluid CSF-Gluc	< 18 y ≥ 18 y	3.33–4.44 2.22–3.89	mmol/L	2 2
CSF-Gluc/S,P-Gluc	≥ 18 y	~0.6		13
Glycated hemoglobin (B-HbA1c)	All age groups	4.8–5.9 29–42	% of total Hb mmol/mol	2
Glucose tolerance test (GTT): • Glucose in serum/plasma, fasting (fS,fP-Gluc 0 min) • Glucose in serum/plasma, 120 min after oral administration of glucose (S,P-Gluc 120 min)	All age groups	<u>Normal:</u> 0 min ≤ 6.0 120 min < 7.8 <u>Diabetes:</u> 0 min ≥ 7.0 120 min ≥ 11.1 <u>Impaired glucose tolerance (IGT):</u> 0 min < 7.0 120 min 7.8–11.0 <u>Impaired fasting glucose (IFG):</u> 0 min 6.1–6.9 120 min < 7.8	mmol/L	15
Glucose-6-phosphate dehydrogenase (RBC-G6PD/B-Hb)	≥ 18 y	8.0–14.5	U/gHb	33
Haloperidol (P-Haloperidol)	All age groups	Therapeutic range 1–10 Toxic > 15	µg/L	11
Acid-base balance (aB-ABB)				5

pH (aB-pH)	≥ 1 d	7.35–7.45		
Oxygen, partial pressure (aB-pO ₂)	≥ 1 d	83–108	mmHg	
Carbon dioxide, partial pressure (aB-pCO ₂)	≥ 18 y	M 35–48 F 32–45	mmHg	
Bicarbonate (aB-HCO ₃)	≥ 18 y	M 24–31 F 22–31	mmol/L	35
Base excess (aB-BE)	≥ 18 y	M (-2.7) -(+2.5) F (-3.4)-(+1.4)		
Haptoglobin (S,P-Hapto)	< 15 d	< 0.1	g/L	5
	15 d – < 1 y	0.1–2.2		
	1 y – < 12 y	0.1–1.6		
	12 y – < 18 y	0.1–1.8		
	≥ 18 y	0.3–2.0		2
Hemoglobin in plasma (P-Hb)	≥ 18 y	< 100	mg/L	5
Holo transcobalamin (S-HoloTC)	20 y – < 80 y	37.5–188	pmol/L	2
Homocysteine (S,P-Hcy)	5 d – < 1 y	< 10.0	µmol/L	44
	1 y – < 7 y	< 7.6		
	7 y – < 12 y	< 8.4		
	12 y – < 15 y	< 10.4		
	15 y – < 19 y	N < 11.9 M < 13.4		
	≥ 19 y	< 12.0		2
5-hydroxyindoleacetic acid in 24h urine (dU-5-HIAA)	≥ 18 y	2–8	mg/d	9
Monoclonal immunoglobulines in serum (S-Monclon-Ig)	All age groups	Normal finding is negative for monoclonal immunoglobulines		
Monoclonal immunoglobulines in urine (U-Monclon-Ig)	All age groups	Normal finding is negative for monoclonal immunoglobulines		
Immunoglobulin A (S,P-IgA)	< 1 y	3.2–12.0	g/L	44
	1 y – < 3 y	1.5–6.3		
	3 y – < 6 y	3.2–9.9		
	6 y – < 14 y	5.0–11.7		
	14 y – < 19 y	6.0–13.1		
	≥ 19 y	0.70–4.00		2
Immunoglobulin G (S,P-IgG)	< 15 d	3.2–12.0	g/L	44
	15 d – < 1 y	1.5–6.3		
	1 y – < 4 y	3.2–9.9		
	4 y – < 10 y	5.0–11.7		
	10 y – < 19 y	6.0–13.1		
	≥ 19 y	7.00–16.00		2
Immunoglobulin G in cerebrospinal fluid (CSF-IgG)	≥ 18 y	10–30	mg/L	2
Immunoglobulin G index (CSF-S-IgG-ind)	≥ 18 y	< 0.6		7
Immunoglobulin M (S,P-IgM)	< 15 d	< 0.3	g/L	44
	15 d – < 13 w	0.1–0.7		
	13 w – < 1 y	0.1–0.8		
	1 y – < 19 y	M 0.4–1.4 N 0.4–1.8		
	≥ 19 y	0.40–2.30		2
Immunoglobulin free light chains:				
• kappa free light chains (S,P-IgKappa free)	≥ 18 y	2.37–20.73	mg/L	2
• lambda free light chains		4.23–27.69		

(S,P-IgLambda free) • ratio: kappa free light chains/lambda free light chains		0.22–1.74																																																						
Insulin (S,P-Ins)	3 y – < 3.5 y 3.5 y – < 4 y 4 y – < 4.5 y 4.5 y – < 5 y 5 y – < 5.5 y 5.5 y – < 6 y 6 y – < 6.5 y 6.5 y – < 7 y 7 y – < 7.5 y 7.5 y – < 8 y 8 y – < 8.5 y 8.5 y – < 9 y 9 y – < 9.5 y 9.5 y – < 10 y 10 y – < 10.5 y 10.5 y – < 11 y ≥ 18 y	<table border="1"> <thead> <tr> <th>Women</th> <th>Men</th> </tr> </thead> <tbody> <tr><td>0.5–8.4</td><td>0.4–7.0</td></tr> <tr><td>0.6–8.7</td><td>0.4–7.7</td></tr> <tr><td>0.6–9.0</td><td>0.5–8.3</td></tr> <tr><td>0.7–9.3</td><td>0.6–8.8</td></tr> <tr><td>0.8–9.6</td><td>0.7–9.2</td></tr> <tr><td>0.9–9.7</td><td>0.8–9.5</td></tr> <tr><td>1.0–9.8</td><td>0.9–9.8</td></tr> <tr><td>1.1–10.0</td><td>1.0–10.0</td></tr> <tr><td>1.3–10.3</td><td>1.1–10.2</td></tr> <tr><td>1.4–10.8</td><td>1.2–10.5</td></tr> <tr><td>1.6–11.4</td><td>1.3–10.9</td></tr> <tr><td>1.9–12.2</td><td>1.5–11.4</td></tr> <tr><td>2.2–13.1</td><td>1.7–12.1</td></tr> <tr><td>2.5–14.1</td><td>1.9–12.9</td></tr> <tr><td>2.8–15.1</td><td>2.1–13.5</td></tr> <tr><td>3.2–16.1</td><td>2.4–14.2</td></tr> <tr><td>2.6–24.9</td><td></td></tr> </tbody> </table>	Women	Men	0.5–8.4	0.4–7.0	0.6–8.7	0.4–7.7	0.6–9.0	0.5–8.3	0.7–9.3	0.6–8.8	0.8–9.6	0.7–9.2	0.9–9.7	0.8–9.5	1.0–9.8	0.9–9.8	1.1–10.0	1.0–10.0	1.3–10.3	1.1–10.2	1.4–10.8	1.2–10.5	1.6–11.4	1.3–10.9	1.9–12.2	1.5–11.4	2.2–13.1	1.7–12.1	2.5–14.1	1.9–12.9	2.8–15.1	2.1–13.5	3.2–16.1	2.4–14.2	2.6–24.9		mU/L	26																
Women	Men																																																							
0.5–8.4	0.4–7.0																																																							
0.6–8.7	0.4–7.7																																																							
0.6–9.0	0.5–8.3																																																							
0.7–9.3	0.6–8.8																																																							
0.8–9.6	0.7–9.2																																																							
0.9–9.7	0.8–9.5																																																							
1.0–9.8	0.9–9.8																																																							
1.1–10.0	1.0–10.0																																																							
1.3–10.3	1.1–10.2																																																							
1.4–10.8	1.2–10.5																																																							
1.6–11.4	1.3–10.9																																																							
1.9–12.2	1.5–11.4																																																							
2.2–13.1	1.7–12.1																																																							
2.5–14.1	1.9–12.9																																																							
2.8–15.1	2.1–13.5																																																							
3.2–16.1	2.4–14.2																																																							
2.6–24.9																																																								
Insulin-like growth factor 1 (S,P-IGF-1)	< 2y 2 y 3 y 4 y 5 y 6 y 7 y 8 y 9 y 10 y 11 y 12 y 13 y 14 y 15 y – < 18 y 18 y – < 22 y 22 y – < 26 y 26 y – < 31 y 31 y – < 36 y 36 y – < 41 y 41 y – < 46 y 46 y – < 51 y 51 y – < 56 y 56 y – < 62 y ≥ 62 y	<table border="1"> <thead> <tr> <th>Women</th> <th>Men</th> </tr> </thead> <tbody> <tr><td>13.8–104</td><td>11.8–96.4</td></tr> <tr><td>26.1–128</td><td>13.9–104</td></tr> <tr><td>34.2–155</td><td>18.9–116</td></tr> <tr><td>43.2–185</td><td>26.8–134</td></tr> <tr><td>53.0–216</td><td>36.6–156</td></tr> <tr><td>63.6–250</td><td>47.1–184</td></tr> <tr><td>75.0–286</td><td>57.5–216</td></tr> <tr><td>87.3–324</td><td>67.5–254</td></tr> <tr><td>99.9–363</td><td>76.9–296</td></tr> <tr><td>112–398</td><td>85.7–343</td></tr> <tr><td>123–427</td><td>93.9–392</td></tr> <tr><td>132–451</td><td>101–434</td></tr> <tr><td>140–468</td><td>108–467</td></tr> <tr><td>146–480</td><td>115–489</td></tr> <tr><td>151–485</td><td>120–503</td></tr> <tr><td>148–466</td><td>132–476</td></tr> <tr><td>130–392</td><td>132–370</td></tr> <tr><td>112–329</td><td>120–295</td></tr> <tr><td>100–271</td><td>109–253</td></tr> <tr><td>91.4–238</td><td>98.5–239</td></tr> <tr><td>83.3–225</td><td>88.5–226</td></tr> <tr><td>75.7–219</td><td>78.8–214</td></tr> <tr><td>68.6–214</td><td>68.9–203</td></tr> <tr><td>60.7–201</td><td>60.0–195</td></tr> <tr><td>55.1–179</td><td>49.6–189</td></tr> </tbody> </table>	Women	Men	13.8–104	11.8–96.4	26.1–128	13.9–104	34.2–155	18.9–116	43.2–185	26.8–134	53.0–216	36.6–156	63.6–250	47.1–184	75.0–286	57.5–216	87.3–324	67.5–254	99.9–363	76.9–296	112–398	85.7–343	123–427	93.9–392	132–451	101–434	140–468	108–467	146–480	115–489	151–485	120–503	148–466	132–476	130–392	132–370	112–329	120–295	100–271	109–253	91.4–238	98.5–239	83.3–225	88.5–226	75.7–219	78.8–214	68.6–214	68.9–203	60.7–201	60.0–195	55.1–179	49.6–189	µg/L	2
Women	Men																																																							
13.8–104	11.8–96.4																																																							
26.1–128	13.9–104																																																							
34.2–155	18.9–116																																																							
43.2–185	26.8–134																																																							
53.0–216	36.6–156																																																							
63.6–250	47.1–184																																																							
75.0–286	57.5–216																																																							
87.3–324	67.5–254																																																							
99.9–363	76.9–296																																																							
112–398	85.7–343																																																							
123–427	93.9–392																																																							
132–451	101–434																																																							
140–468	108–467																																																							
146–480	115–489																																																							
151–485	120–503																																																							
148–466	132–476																																																							
130–392	132–370																																																							
112–329	120–295																																																							
100–271	109–253																																																							
91.4–238	98.5–239																																																							
83.3–225	88.5–226																																																							
75.7–219	78.8–214																																																							
68.6–214	68.9–203																																																							
60.7–201	60.0–195																																																							
55.1–179	49.6–189																																																							
Interleukin 6 (S-IL-6)	≥ 18 y	< 7	ng/L	2																																																				
Isoniasiid (S,P-Isoniazid)		Peak concentration 3–6 (300 mg/day) 9–15 (900 mg twice a weak)	mg/L	49																																																				

Potassium (S,P-K)	1 d – < 8 d 8 d – < 1 m 1 m – < 7 m 7 m – < 1 y 1y – < 18 y ≥ 18 y	3.2–5.5 3.4–6.0 3.5–5.6 3.5–6.1 3.3–4.6 3.4–4.8	mmol/L	1 5
Potassium in urine dU-K	6 y – < 10 y 10 y – < 15 y ≥ 15 y	M 17–54 F 8–37 M 22–57 F 18–58 25–125	mmol/d	5 1, 2, 5
U-K (first morning urine)	≥ 18 y	20–80	mmol/L	1
Calprotectin in stool (St-Calpro)	6 m – < 2 y 2 y – < 4 y ≥ 4 y	< 250 < 100 ≤ 50	µg/g	47
Calcitonin (S,P-CT)	< 3 m 3 m – < 6 m 6 m – < 9 m 9 m – < 18 m 18 m – < 3 y 3 y – < 17 y ≥ 17 y	≤ 10 ≤ 8.0 ≤ 6.4 ≤ 5.0 ≤ 3.0 ≤ 2.0 M ≤ 2.78 N ≤ 1.87	pmol/L	46 2
Calcium (S,P-Ca)	< 11 d 11 d – < 3 y 3 y – < 13 y 13 y – < 18 y 18 y – < 60 y 60 y – < 90 y ≥ 90 y	1.90–2.60 2.25–2.75 2.20–2.70 2.10–2.55 2.15–2.50 2.20–2.55 2.05–2.40	mmol/L	1 2
Calcium (ionized) (S,P-iCa)	≥ 18 y	1.16–1.32	mmol/L	1
Calcium in urine dU-Ca	< 18 y ≥ 18 y	< 0.15 2.5–7.5	mmol/kg/d mmol/d	1 2
U-Ca/U-Crea	6 m – < 1 y 1 y – < 2 y 2 y – < 3 y 3 y – < 5 y 5 y – < 7 y 7 y – < 18y	0.09–2.2 0.07–1.5 0.06–1.4 0.05–1.1 0.04–0.8 0.04–0.7	mol/mol	4
Cannabinoids in urine (U-THC)	All age groups	negative		
Carbamazepine (S,P-Carba)	All age groups	Therapeutic range 4–12 Toxic > 15	mg/L	2, 5
Carboxyhemoglobin (B-COHB)	≥ 18 y	0.5–1.5	% of total Hb	5
Carcinoembryonic antigen (S,P-CEA)	20 y – < 70 y	Non-smokers < 3.8 Smokers < 5.5	µg/L	1, 2
Carbohydrate antigen 15-3 (S,P-CA 15-3)	≥ 18 y	≤ 25	KU/L	1, 2
Carbohydrate antigen 19-9 (S,P-CA 19-9)	≥ 18 y	< 27	KU/L	1, 2
Carbohydrate antigen 72-4 (S,P-CA 72-4)	≥ 18 y	< 6.9	KU/L	2
Carbohydrate antigen 125 (S,P-CA 125)	≥ 18 y	F < 35	KU/L	1, 2
HE4 (S,P-HE4)	Premenopausal Postmenopausal	< 70 < 140	pmol/L	2

ROMA value	Premenopausal Postmenopausal	< 11.4 low risk of finding epithelial ovarian cancer < 29.9 low risk of finding epithelial ovarian cancer	%	2
S-100 (S-S-100)	≥ 18 y	≤ 0.105	µg/L	1, 2
Growth hormone (S,P-GH)	< 11 y 11 y – < 18 y ≥ 18 y	Men 0.280–18.87 Women 0.360–23.37 0.231–32.40 0.369–24.15 <0.09–7.41 0.378–29.64	mU/L	2
Clarithromysin (S,P-Clarithromycin)		Peak concentration >2	mg/L	49
Chloride (S,P-Cl)	1 d – < 7 m 7 m – < 1 y 1 y – < 18 y ≥ 18 y	97–108 97–106 97–107 98–107	mmol/L	1 1, 2
Chloride in sweat (Sw-Cl)		normal < 30 borderline 30–60 cystic fibrosis > 60	mmol/L	28
Chloride in urine dU-Cl	< 1 y 1 y – < 6 y 6 y – < 10 y 10 y – < 15 y ≥ 15 y ≥ 18 y	2–10 15–40 M 36–110 F 18–74 M 64–176 F 36–173 110–250 46–168	mmol/d	5 1, 2, 5 1
U-Cl (first morning urine)			mmol/L	
Cholesterol (S,P-Chol)	1 d – < 1 m 1 m – < 6 m 6 m – < 1 y 1 y – < 4 y 4 y – < 7 y 7 y – < 10 y 10 y – < 13 y 13 y – < 16 y 16 y – < 18 y ≥18 y recommended	M 1.40–3.90 F 1.60–4.01 M 2.09–3.80 F 1.60–3.65 M 1.97–4.63 F 1.97–5.59 M 2.20–4.71 F 2.79–4.99 M 2.84–5.61 F 2.74–4.99 M 2.84–5.46 F 2.69–5.43 M 2.72–5.77 F 2.72–5.64 M 2.35–5.28 F 2.79–5.30 M 2.12–4.97 F 2.38–6.05 < 5.0	mmol/L	1 16
HDL-Cholesterol (S,P-HDL-Chol)		M > 1.0 N > 1.2	mmol/L	16
Non-HDL-Cholesterol (S,P-non-HDL-Chol)	≥18 y recommended	< 3.9	mmol/L	16
LDL-Cholesterol (S,P-LDL-Chol)	≥18 y recommended	< 3	mmol/L	16
Complement component 3 (S,P-C3)	< 15 d 15 d – < 1y 1 y – < 19 y	0.57–1.29 0.58–1.69 0.9–1.61	g/L	44

	≥ 19 y	0.9–1.8		1, 2
Complement component 4 (S,P-C4)	< 1 y	0.07–0.31	g/L	44
	1 y – < 19 y	0.13–0.38		
	≥ 19 y	0.1–0.4		
Cholinesterase (S,P-ChE)	< 16 y	5320–12920	U/L	1, 2
	≥ 16 y	M 5320–12920		
	16 y - < 40 y	F 4260–11250		
	≥ 40 y	F 5320–12920		
	Pregnant woman, or woman using oral contraceptives (18 y – < 42 y)	F 3650–9120		
	Dibucain number	≥ 73		
	Phenotype UU	57–72		
Phenotype UA	≤ 72			
Phenotype AA	<57			
Risk of scolion-apnoe				
Chorionic gonadotropin (intact + β subunit) (S,P-hCG intact + β subunit)	≥ 18 y	M < 2.0	U/L	2
		F nonpregnant ≤ 1.0		
		postmenopausal ≤ 7.0		
		During pregnancy		
		3. weak 5,8–71,2		
		4. weak 9,5–750		
		5. weak 217–7138		
		6. weak 158–31795		
		7. weak 3697–163563		
		8. weak 32065–149571		
		9. weak 63803–151410		
		10. weak 46509–186977		
		12. weak 27832–210612		
		14. weak 13950–62530		
		15. weak 12039–70971		
		16. weak 9040–56451		
		17. weak 8175–55868		
		18. weak 8099–58176		
		Chorionic gonadotropin, free β subunit (S-fβ-hCG)		
Cortisol (S,P-Cort)	< 1 k	15–396	nmol/L	44
	1 k – < 1 a	18–552		
	1 a – < 12 a	66–410		
	12 a – < 19 a	100–480		
	≥ 19 a	6–10 a.m 133–537		
		4–8 p.m 68.2–327		

Cortisol in saliva (Sal-Cort)	≥ 18 y	6.00 – 10.00 < 24.1 16.00 – 20.00 < 9.65 23.30 – 00.30 < 11.3	nmol/L	2
Creatine kinase (S,P-CK)	< 2 d 2 d – < 6 d 6 d – < 7 m 7 m – < 1 y 1 y – < 4 y 4 y – < 7 y 7 y – < 13 y 13 y – < 18 y ≥ 18 y	< 712 < 652 < 295 < 203 < 228 < 149 M < 247 F < 154 M < 270 F < 123 M < 308 F < 192	U/L	1
				2
Creatine kinase, MB isoenzyme, mass (S,P-CK-MBm)	≥ 18 y	M < 6.22 F < 4.88	µg/L	2
Creatinine (S,P-Crea)	Preterm neonates < 2 m 2 m – < 1 y 1 y – < 3 y 3 y – < 5 y 5 y – < 7 y 7 y – < 9 y 9 y – < 11 y 11 y – < 13 y 13 y – < 15 y ≥ 15 y	29–87 27–77 14–34 15–31 23–37 25–42 30–47 29–56 39–60 40–68 M 59–104 F 45–84	µmol/L	2
Creatinine in urine U-Crea (first morning urine) dU-Crea	≥ 18 y	M 3.5–24.6 F 2.6–20.0	mmol/L	2
	3 y – < 9 y 9 y – < 13 y 13 y – < 18 y	0.97–6.0 1.5–12.5 2.6–16.5	mmol/d	3
	≥ 18 y	M 9.0–19.0 F 6.0–13.0		2
Creatinine clearance	5 d – < 8 d 1 m – < 3 m 3 m – < 1 y 3 y – < 14 y ≥ 18 y	> 38 > 54 > 64 > 120 66–143	mL/min/1,73m ²	1 (2004)
			mL/min/1,73m ²	1 (2004)
Estimated glomerular filtration rate (eGFR (Crea, CKD-EPI))	≥ 18 y	≥ 90	mL/min/1,73m ²	12
Cryoglobulins (S-Cryo)	≥ 18 y	< 50	mg/L	38
Uric acid (S,P-UA)	1 d – < 1 m 1 m – < 1 y 1 y – < 4 y 4 y – < 7 y 7 y – < 10 y 10 y – < 13 y 13 y – < 16 y	M 71–230 F 59–271 M 71–330 F 65–319 M 124–330 F 106–295 M 106–325 F 118–301 M 106–319 F 106–325 M 130–342 F 148–348 M 183–413 F 130–378	µmol/L	1

	16 y – < 19 y ≥ 19 y	M 124–448 F 142–389 M 202–417 F 143–339 Target level in case of podagra therapy < 360		2
Uric acid in urine dU-UA U-UA (first morning urine)	≥ 18 y ≥ 18 y	1.2–5.9 2.2–5.5	mmol/d mmol/L	1, 2 1, 2
Lactate dehydrogenase (S,P-LDH)	< 15 p 15 p – < 1a 1a – < 10 a 10 a – < 15 a 15 a – < 19 a ≥ 19 a	< 1128 < 424 < 305 < 260 < 240 < 250	U/L	44 2
Lactate dehydrogenase in pleural fluid (PirF-LDH) expressed by ratio PirF-LDH/S,P-LDH	All age groups	Transudate < 0.6 Exudate > 0.6		13, 1
Lactate vP-Lac aP-Lac	≥ 18 y ≥ 18 y	< 2.2 < 1.6	mmol/L	1, 2
Lactate in cerebrospinal fluid (CSF-Lac)	< 3 d 3 d – < 11 d 11 d – < 18 y ≥ 18 y	1.1–6.7 1.1–4.4 1.1–2.8 1.1–2.4	mmol/L	1, 2
Lactose tolerance test (LTT)	All age groups	A rise in the blood glucose concentration > 1.1 mmol/L indicates the absence of lactase deficiency		20
Lamotrigine (S,P-Lamotr)	All age groups	Therapeutic 3.0–15 Toxic > 20	mg/L	9
Leflunomiid (P-Leflunomide)	All age groups	Therapeutic range is not uniquely determined. Before conception level must be < 0.02	mg/L	10
Levetiracetam (S,P-Levetir)	All age groups	12–46	mg/L	9
Lithium (S-Li)	≥ 18 y	Therapeutic 0.6–1.2 Toxic > 2	mmol/L	2
Linezolid (S,P-Linezolid)		Peak concentration 12–26	mg/L	49
Lipase (S,P-Lip)	< 1 m 1 m – < 13 y 16 y – < 18 y ≥ 18 y	< 34 < 31 < 55 13–60	U/L	1 2
Luteinizing hormone (S,P-LH)	< 6 m 6 m – < 11 y 11 y – < 14 y 14 y – < 18 y ≥ 18 y	M < 6.2 F < 8.2 M < 1.3 F < 1.3 M < 2.0 F < 10.0 M 1.3–8.4 F 0.4–25 M 1.7–8.6 F follic.phase 2,4–12,6 Ovul 14–95,6 lut.phase 1.0–11.4	U/L	44 1, 2

		postmenop 7.7–58.5		
Magnesium (S,P-Mg)	2 d – < 5 d 5 m – < 6 y 6 y – < 12 y 12 y – < 20 y 20 y – < 60 y 60 y – < 90 y ≥ 90 y	0.62–0.91 0.70–0.95 0.70–0.86 0.70–0.91 0.66–1.07 0.66–0.99 0.70–0.95	mmol/L	2, 5
Magnesium in urine dU-Mg U-Mg	≥ 18 y ≥ 18 y	3.0–5.0 1.7–5.7	mmol/d mmol/L	2 1 (2004)
Methadone in urine (U-Mtd)	All age groups	negative		
Metamphetamines in urine (U-Met)	All age groups	negative		
Methemoglobin (B-MetHb)	≥ 18 y	< 0.6	% of total Hb	35
Meropenem (P-Meropenem)	All age groups	Therapeutic range is not uniquely determined, effect is assessed using minimal inhibitory concentration (MIC)	mg/L	21
Methotrexate (S,P-MTX)	All age groups	Therapeutic range depends on dose of MTX and specimen collecting time	µmol/L	39
Moxifloxacin (S,P-Moxifloxacin)		Peak concentration 3–5	mg/L	49
Mycophenolic acid (P-MPA)	All age groups	Depends on the type of transplantate and concomitant administration of some other drugs		2
Myoglobin (S,P-Myogl)	≥ 18 y	M 28–72 F 25–58	µg/L	1, 2
Sodium (S,P-Na)	< 8 d 8 d – < 2 m 2 m – < 7 m 7 m – < 1 y 1 y – < 18 y ≥ 18 y	131–144 132–142 132–140 131–140 132–141 136–145	mmol/L	1 2
Sodium in urine dU-Na U-Na (first morning urine)	6 y – < 10 y 10 y – < 15 y ≥ 15 y ≥ 18 y	M 41–115 F 20–69 M 63–177 F 48–168 M 40–220 F 27–287 54–190	mmol/d mmol/L	5 1
Neuron specific enolase (S-NSE)	≥ 18 y	< 16.3	µg/L	2
Oxcarbazepin (S,P-Oxcarb)	All age groups	MHD conc 3.6–35	mg/L	9
Olanzapine (P-Olanzapine)	All age groups	Therapeutic 20–80 Toxic > 100	µg/L	11
Oligoclonal immunoglobulin G in cerebrospinal fluid (CSF-IgG-oligo)	All age groups	Normal finding is negative		

Opiates in urine (U-Mop)	All age groups	negative		
Osmolality (S-Osmol)	18 y – < 61 y ≥ 61 y	275–295 280–300	mosm/kgH ₂ O	1
Osmolality in urine (U-Osmol)	≥ 18 y	400–800	mosm/kgH ₂ O	1
Parathyroid hormone (S-PTH)	< 1 k 1 k – < 1 a 1 a – < 11 a 11 a – < 19 a ≥ 19 a	0,7–6,3 0,9–6,5 1,2–6,3 1,6–7,2 M, F 1.6–6.9	pmol/L	44 1, 2
Paracetamol (S,P-Paracet)	≥ 18 y	Therapeutic range 10–30 Toxic > 200 (4 h after administration) > 100 (8 h after administration) > 50 (12 h after administration)	mg/L	2
Piperacillin (P-Piperacillin)	All age groups	Therapeutic range is not uniquely determined, effect is assessed using minimal inhibitory concentration (MIC)	mg/L	21
Posaconazole (P-Posaconazole)	All age groups	Therapeutic range is not uniquely determined, in case of prophylactic therapy suggested concentration is > 0,7	mg/L	18
Porphobilinogen in urine U-PBG U-PBG/U-Crea	≥ 18 y ≥ 18 y	< 8.84 < 1.5	µmol/L mmol/mol	31 6
Porphyrines in urine U-Porph U-Porph/U-Crea	≥ 18 a ≥ 18 a	20–320 < 38	nmol/L µmol/mol	31 36
Prealbumin (S-PreAlb)	< 15 d 15 d – < 1 y 1 y – < 5 y 5 y – < 13 y 13 y – < 16 y 16 y – < 19 y ≥ 19 y	< 0.11 0.04–0.24 0.11–0.23 0.13–0.26 0.17–0.31 M 0.2–0.35 N 0.16–0.33 0.20–0.40	g/L	1 1, 2
Progesterone (S,P-Prog)	F 1 m – < 12 y F 12 y – < 19 y M 1 m – < 19 y ≥ 19 y	< 3 < 38 < 3 M 0.7–4.3 F follic.phase 0.6–4.7 ovulat.phase 2.4–9.4 lut.phase 5.3–86 postmenop. 0.3–2.5	nmol/L	44 1, 2
Procalcitonin (S,P-PCT)	< 6 h 6 h – < 12 h 12 h – < 18 h	< 2 < 8 < 15	µg/L	50

	18 h – < 30 h 30 h – < 36 h 36 h – < 42 h 42 h – < 48 h ≥ 3 d	< 21 < 15 < 8 < 2 < 0.05 Value > 2 is indicative for sepsis		2
Prolactin (S,P-Prol)	1 m – < 1 y 1 y – < 19 y ≥ 19 y	110–1274 64–532 M 86–324 F 102–496	mU/L	44 2
Prostate-specific antigen (S,P-PSA)	< 40 y 40 y – < 50 y 50 y – < 60 y 60 y – < 70 y ≥ 70 y	< 1.4 < 2.0 < 3.1 < 4.1 < 4.4	µg/L	1, 2
Free prostate-specific antigen (S,P-fPSA%)	All age groups	The risk of prostate cancer increases if fPSA% is < 15–25%	%	2
Pyrasinamid (S,P-Pyrasinamid)		Peak concentration 20–60 (25–35 mg/kg day) 60–90 (50 mg/kg day)	mg/L	49
Pregnancy associated protein A (S-PAPP-A)		Result is considered in complex of I trimester pregnancy screening	U/L	
Iron (S,P-Fe)	< 14 y 14 y – < 19 y ≥ 19 y	M, F 5.0–25.0 M 8.0–31.0 F 6.0–31.0 M 11.0–28.0 F 6.6–26.0	µmol/L	44 1
Renin (P-Renin)	14 d – < 4 m 4 m – < 1 y 1 y – < 3 y 3 y – < 5 y 5 y – < 7 y 7 y – < 11 y 11 y – < 15 y 15 y – < 18 y ≥ 18 y	11.2–147.9 supine 17.4–173.8 supine 21.4–102.3 supine 19.5–123.0 supine 20.4–128.8 supine 14.8–102.3 supine 13.8–104.7 supine 13.8–72.4 supine 5.3–99.1 upright	mU/L	43 42
Rheumatoid factor (S,P-RF)	≥ 18 y	< 14	kU/L	1, 2
Risperidone and 9-hydroksyrisperidoyne (P-Risperidone+9-hydroksyrisperidone)	All age groups	Therapeutic 20–60 Toxic > 120	µg/L	11
Salicylates (S,P-Salic)	≥ 18 y	Therapeutic range: antipyretic, analgetic 30–100 anti-inflammatory 150–300 Toxic > 300 lethal > 600	mg/L	2
Sirolimus (B-Sirolimus)	All age groups	Therapeutic range depends on indications of Sirolimus administration and	µg/L	2

		specimen collection time		
Sex hormone binding protein (S,P-SHBG)	< 1 k 1 k – < 13 a 13 a – < 15 a 15 a – < 19 a 19 a – < 50 a ≥ 50 y	> 16 > 37,5 21,1–152 M 13,6–62 N 21,6– 127,0 M 18.3–54.1 F 32.4– 128	nmol/L	44
Free androgen index (FAI)	20 y – < 50 y ≥ 50 y	M 20.6–76.7 F 27.1– 128 M 35.0–92.6 F 0.297–5.62 M 24.30–72.1 F 0.187–3.63	%	2
Carbohydrate deficient transferrin (IFCC) S-CDT (IFCC)	≥ 18 y	≤ 1.7% excessive alcohol consumption in the past two weeks is unlikely 1.8–2.0% borderline result, interpretation in relation to excessive alcohol consumption is not possible ≥ 2.1% the result indicates excessive alcohol consumption in the past two weeks	%	41
Tacrolimus (B-Tacro)	All age groups	Therapeutic range depends on indications of Tacrolimus administration and specimen collection time	µg/L	2
Theophylline (S,P-Theoph)	All age groups (except newborns) ≥ 18 y	Therapeutic range for bronchodilatation: 10–20 Treatment of neonatal apnoe: 6–13 Toxic: > 20	mg/L	1, 2 5 5
Testosteron (S,P-Testo)	< 6 m 6 m – < 11 y 11 y – < 19 y 11 y – < 15 y 15 y – < 19 y 19 y – < 50 y ≥ 50 y	M 0.2–19 F < 12 M, F < 0.10 F < 1.8 M < 20 M 1.7–24 M 8.64–29.0 F 0.29–1.67 M 6.68–25.7 F 0.10–1.42	nmol/L	44
Free testosteron (S,P-fTesto)	18 y – < 50 y	M > 0.220	nmol/L	19

calc)	≥ 50 y	M > 0.180		
Thiopurine methyltransferase in erythrocytes (RBC-TPMT) Performer : Departement of Pharmacology, Tartu University	≥ 18 y	Concentration of 6-methylmercaptapurine after 60 min of incubation 59–110	ng/mL(RBC)/h	
Transferrin (S,P-Transf)	< 9 w 9 w – < 1 y 1 y – < 19 y ≥ 19 y	1.11–2.43 1.15–3.52 2.38–3.66 2.0–3.6	g/L	44 2
Transferrin saturation	≥ 18 y	16–45	%	1
Soluble transferrin receptor (S,P-Transf-sR)	9 m – < 1 y 18 y – < 83 y	4.1–7.7 1.71–4.13	mg/L	22 2
Triglycerides (S,P-Trigl)	Preterm neonates ≥ 18 y	< 0.7 recommended < 1.7 fasting < 2.0 nonfasting	mmol/L	1 16, 2
Free triiodthyronine (S,P-ft3)	< 6 d 6 d – < 4 m 4 m – < 1 y 1 y – < 7 y 7 y – < 12 y 12 y – < 21 y ≥ 21 y Pregnancy	2.65–9.68 3.00–9.28 3.30–8.95 3.69–8.46 3.88–8.02 3.93–7.70 3.10–6.80 I trim 3.8–6.0 II trim 3.2–5.5 III trim 3.1–5.0	pmol/L	1 1, 2 1
Tricyclic antidepressants in urine (U-TCA)	All age groups	negative		
Troponin T (high sensitivity) (S,P-cTnT-hs)	≥ 18 y	< 14 ≥ 14 myocardial necrosis	ng/L	2, 29
Ceruloplasmin (S,P-Cer)	< 2 m 2 m – < 6 m 6 m – < 1 y 1 y – < 8 y 8 y – < 14 y 14 y – < 19 y ≥ 19 y	0.07–0.24 0.13–0.33 0.14–0.39 0.22–0.43 0.21–0.40 M 0.17–0.35 N 0.21–0.43 M 0.15–0.30 F 0.16–0.45	g/L	5, 44 2
Zinc (S-Zn)	≥ 18 y	M 11.1–19.5 F 10.7–17.5	µmol/L	14
Antibodies to cyclic citrullinated peptide (S,P-CCP IgG)	All age groups	< 17	kU/L	2
Cycloserine (S,P-Cycloserine)	All age groups	Peak concentration: Therapeutic 20–35 Toxic > 35	mg/L	49

Cyclosporin A (B-CyA)	All age groups	Therapeutic range depends on indications of CyA administration and specimen collection time	µg/L	2
Cystatin C (S,P-CysC)	< 1 m 1 m – < 1 y 1 y – < 2 y 2 y – < 3 y 3 y – < 5 y 5 y – < 6 y 6 y – < 9 y 9 y – < 10 y 10 y – < 11 y 11 y – < 12 y 12 y – < 13 y 13 y – < 14 y 14 y – < 15 y 15 y – < 16 y 16 y – < 17 y 17 y – < 18 y 18 y – < 78 y	1.1–2.2 0.5–1.4 M 0.74–1.22 F 0.74–1.20 M 0.67–1.10 F 0.67–1.08 M 0.65–1.06 F 0.64–1.04 M 0.65–1.07 F 0.66–1.06 M 0.65–1.09 F 0.67–1.08 M 0.66–1.10 F 0.68–1.09 M 0.66–1.11 F 0.68–1.11 M 0.67–1.13 F 0.69–1.14 M 0.69–1.17 F 0.68–1.16 M 0.72–1.22 F 0.66–1.14 M 0.74–1.24 F 0.64–1.11 M 0.74–1.23 F 0.63–1.09 M 0.73–1.20 F 0.62–1.07 M 0.71–1.15 F 0.61–1.05 0.61–0.95	mg/L	1 30 2
Estimated glomerular filtration rate (eGFR (CysC, CKD-EPI))	≥ 18 y	≥ 90	mL/min/1,73m ²	12
Thyroglobulin (S,P-TG)	< 6 d 6 d – < 4 m 4 m – < 1 y 1 y – < 7 y 7 y – < 12 y 12 y – < 21 y ≥ 21 y	25–307 20–228 18–125 9.0–67 5.1–43 2.6–36 1.4–78	µg/L	1 2
Antibodies to thyroglobulin (S-TG IgG)	< 6 d 6 d – < 4 m 4 m – < 1 y 1 y – < 7 y 7 y – < 12 y 12 y – < 21 y ≥ 21 y	< 134 < 146 < 130 < 38 < 37 < 64 < 115	KU/L	1 1, 2
Antibodies to thyroid peroxidase (S,P-TPO IgG)	< 6 d 6 d – < 4 m	< 117 < 47	KU/L	1

	4 m – < 1 y 1 y – < 7 y 7 y – < 12 y 12 y – < 21 y ≥ 21 y	< 32 < 13 < 18 < 26 < 34		1, 2
Thyroid-stimulating hormone (S,P-TSH)	< 6 d 6 d – < 4 m 4 m – < 1 y 1 y – < 7 y 7 y – < 12 y 12 y – < 21 y ≥ 21 y	0.70–15.2 0.72–11.0 0.73–8.35 0.70–5.97 0.60–4.84 0.51–4.30 0.27–4.2	mU/L	1 1, 2
Thyroid stimulating immunoglobulins (S,P-TSI)	≥ 18 y	< 0.1 > 0.55 cut off for Graves disease	U/L	48
Free thyroxine (S,P-ft4)	< 6 d 6 d – < 4 m 4 m – < 1 y 1 y – < 7 y 7 y – < 12 y 12 y – < 21 y ≥ 21 y Pregnancy	11.0–32.0 11.5–28.3 11.9–25.6 12.3–22.8 12.5–21.5 12.6–21.0 12.0–22.0 I trim 12.1–19.6 II trim 9.6–17.0 III trim 8.4–15.6	pmol/L	1 1, 2 1
Urea (S,P-Urea)	0 d – < 15 d 15 d – < 1 y 1 y – < 10 y 10 y – < 19 y ≥ 19 y	1.1–7.9 1.3–5.8 3.2–7.6 F 2.6–6.5 M 2.6–7.2 < 8.1	mmol/L	44 2
Urea in urine dU-Urea U-Urea (first morning urine)	≥ 18 y ≥ 18 y	428–714 286–595	mmol/d mmol/L	2 2
Protein (S,P-Prot)	< 15 d 15 d – < 1 y 1 y – < 6 y 6 y – < 9 y 9 y – < 19 y ≥ 19 y	51–80 43–69 59–73 62–75 63–78 64–83	g/L	44 1, 2
Protein in cerebrospinal fluid (CSF-Prot)	1 d – < 2 m 2 m – < 4 m 4 m – < 7 m 7 m – < 1 y 1 y – < 3 y 3 y – < 5 y 5 y – < 9 y ≥ 18 y	0.25–0.72 0.20–0.72 0.15–0.50 0.10–0.45 0.10–0.40 0.10–0.38 0.10–0.43 0.15–0.45	g/L	1 2
Protein in pleural fluid (PlrF-Prot) PlrF-Prot/S,P-Prot	All age groups	transudate < 0.5 exudate > 0.5	g/L	13
Protein in urine dU-Prot U-Prot/U-Crea	≥ 18 y ≥ 18 y	< 0.15 < 0.2	g/d mg/mg	1 27

Protein fractions in serum (S-Prot-Fr)	≥ 18 y	albumin 35–52 globulins: alpha 1 1.0–3.0 alpha 2 4.0–8.0 beta 1 4.0–8.0 beta 2 2.0–7.0 gamma 7.0–17.0	g/L	
Protein fractions in urine (U-Prot-Fr)	All age groups	Results will be commented by laboratory doctor	mg/L	1, 2
Valproate (S,P-Valpr)	All age groups	Therapeutic 50–100 Toxic > 150	mg/L	1,2 2
Free valproate (S,P-fValpr)	All age groups	Therapeutic 5–15	mg/L	24
Free valproate % (S-fValpr%)		5–15	%	52
Vancomycin (S,P-Vanco)	≥ 18 y	Pre-dose (trough) concentration: therapeutic range 15–20 AUC/MIC 400–600 (if MIC is ~ 1 mg/L)	mg/L	40 27
Bicarbonate (S,P-HCO ₃)	≥ 18 y	22–29	mmol/L	1, 2
Vitamin A (S,P-Vit A)	< 1 y 1 y – < 7 y 7 y – < 13 y 13 y – < 20 y ≥ 20 y	0.10–0.50 0.20–0.43 0.26–0.49 0.26–0.72 0.30–0.70	mg/L	9
Vitamin B1 (B-Vit B1)	All age groups	33.1–60.7	µg/L	9
Vitamin B6 (B-Vit B6)	All age groups	12.6–45.2	µg/L	9
Vitamin B12 (S,P-Vit B12)	< 1 m 1 m – < 1 y 1 y – < 12 y 12 y – < 19 y ≥ 19 y	138–1377 124–1236 261–1180 199–835 145–569	pmol/L	44 2
Vitamin E (S,P-Vit E)	< 1 y 1 y – < 7 y 7 y – < 13 y 13 y – < 20 y ≥ 20 y	1–8 3–9 4–9 6–10 5–20	mg/L	9
Vitamin D (S,P-Vit D(25-OH))	All age groups	Deficiency < 50	nmol/L	34
Vorikonazol (P-Voricon)	≥ 18 y	1.0–5.5	mg/L	9
Estradiol (S,P-E2)	1 m – < 18 y 1 m – < 10 y 10 y – < 14 y 14 y – < 18 y ≥ 18 y	M < 18 N < 18 N < 250 N 53.6–912 M < 159 F follic.phase 45–854 ovulat.phase 151–1461 lut.phase 82–1251 postmenop. < 183	pmol/L	44 2

References

- 1 W.Heil, R. Koberstein, B.Zawta. Reference Ranges for Adults and Children. Pre-Analytical Considerations. Roche 2008
- 2 Cobas method sheets
- 3 S.J. Soldin, C.Brugnara, E.C.Wong. Pediatric Reference Ranges. AACCC Press 2003
- 4 V.Matos, G.Melle, O.Boulat et al. Urinary phosphate/creatinine, calcium/creatinine and magnesium/creatinine ratios in a healthy pediatric population. The Journal of Pediatrics, volume 131, number 2
- 5 Burtis, Ashwood, Bruns. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics. Elsevier Saunders 2018
- 6 European Porphyria Network (<https://porphyria.eu/>)
- 7 K.Kallion, K.Reimand, A.Orav. Comparison of the IgG index and oligoclonal IgG bands in CSF: finding out optimal cut-off value for IgG index. Abstract No: P 29 at the 31th Nordic Congress in Clinical Chemistry, June 14-18, Helsinki, Finland
- 8 TIAFT reference blood level list of therapeutic and toxic substances, september 2004
- 9 Chromsystems. Method sheets
- 10 Krause M, Amin S, Makol A. Use of DMARDs and biologics during pregnancy and lactation in rheumatoid arthritis: what the rheumatologist needs to know. Ther Adv Musculoskel Dis 2014; 6(5) 169–184
- 11 Hiemke C et al. Consensus Guidelines for Therapeutic Drug Monitoring in Neuropsychopharmacology: Update 2017. Pharmacopsychiatry 2018; 51: 9–62
- 12 KDIGO 2012 Clinical practice guideline for the evaluation and management of chronic kidney disease. Kidney Int 2013; vol 3, issue 1
- 13 Brunzel NA. Fundamentals of Urine and Bodyfluid Analysis. Saunders 2018
- 14 Randox Zinc Colorimetric Method Cat. No. ZN 2341 Reagent Information
- 15 Eesti 2. tüüpi diabeedi juhend 2016
- 16 European guidelines on cardiovascular disease prevention in clinical practice, 2016
- 2019 ESC/EAS Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk
- 17 Bühlmann. ACE kinetic. 2022-10-21
- 18 Ashbee HR et al. Therapeutic drug monitoring of antifungal agents: guidelines from the British Society for Medical Mycology. J Antimicrob Chemother 2014; 69:1162–1176
- 19 Bhasin S, Pencina M et al. Reference Ranges for Testosterone in Men Generated Using Liquid Chromatography Tandem Mass Spectrometry in a Community-Based Sample of Healthy Nonobese Young Men in the Framingham Heart Study and Applied to Three Geographically Distinct Cohorts. J Clin Endocrinol Metab. 2011 Aug; 96(8): 2430–2439
- 20 McPherson RA, Pincus MR. Henry's Clinical Diagnosis and Management by Laboratory Methods. Saunders 2007
- 21 Agency of Medicines. Register of medicinal products. Summary of product characteristic
- 22 Vendt, N. Rauapuudus ja rauapuudusaneemia 9-12 kuu vanustel imikutel Eestis. Doktoriväitekiri 2008
- 23 EV Liiklusseadus, 01.07.2011
- 24 UpToDate. Valproate: drug information
- 25 Nir A., Lindiger A et al. NT-pro-B-type Peptide in Infants and Children: Reference Values Based on Combined Data from Four Studies. Pediatr Cardiol 2009;30:3–8

- 26 Peplies J, Jiménez-Pavón D, Savva SC et al. Percentiles of fasting serum insulin, glucose, HbA1c and HOMA-IR in pre-pubertal normal weight European children from the IDEFICS cohort. *International Journal of Obesity* 2014; 38:39–S47
- 27 Rybak M, Le J. Therapeutic Monitoring of Vancomycin for Serious Methicillin-resistant *Staphylococcus aureus* Infections: A Revised Consensus Guideline and Review by the American Society of Health-system Pharmacists, the Infectious Diseases Society of America, the Pediatric Infectious Diseases Society, and the Society of Infectious Diseases Pharmacists. *Am J Health-Syst Pharm.* 2020;77:835-864
- 28 K.de Boeck, M.Wilschanski, C.Castellani et al. Cystic fibrosis: terminology and diagnostic algorithms. *Thorax* 2006; 61:627-635
- 29 Müokardiinfarkti nelja universaalne definitsioon, *Eesti Arst* 2019; 98(4):235–242
- 30 Ziegelasch N, Vogel M et al. Cystatin C serum levels in healthy children are related to age, gender and pubertal stage. *Pediatric Nephrology* 2019; 34:449–457
- 31 BioRad. ALA/PBG by column test, Sept 2015
- 32 Immulite Compendium, 2007
- 33 Koitsaar L. Kineetilisel ensümaatilisel meetodil määratava glükoos-6-fosfaadi dehüdrogenaasi referentsväärtusvahemiku leidmine Eesit populatsioonile. Tartu Tervishoiu Kõrgkooli bioanalüütiku õppekava lõputöö. Tartu 2023
- 34 Kull M, Kallikorm R, Lember M. D-vitamiin – taasleitud oluline tervisemõjur. *Eesti Arst* 2010; 89(3):185–190
- 35 The Blood Gas Handbook. Radiometer Copenhagen.2000
- 36 Karolinska Universitetslaboratoriet (<https://www.karolinska.se/KUL>)
- 37 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure, 2016
- 38 Le Carrer D. Serum Protein Electrophoresis and immunofixation. Laboratories Sebia, 2005
- 39 Alinity i, method sheet, Abbott
- 40 Rybak M, Lomaestro B et al. Therapeutic monitoring of vancomycin in adult patients: A Consensus Review of the American Society of Health-System Pharmacists, the Infectious Diseases Society of America and the Society of Infectious Diseases Pharmacists. *Am J Health-Syst Pharm.* 2009; 66:82-98
- 41 Sebia, method sheets
- 42 IDS-iSYS method sheets
- 43 Krüger C, Rauh M., Dörr H. Immunoreactive renin concentrations in healthy children from birth to adolescence. *Clinica Chimica Acta* 1998; 274: 15-27
- 44 CALIPER database
- 45 Novanet method sheet
- 46 Castagna G et al. Reference Range of serum calcitonin in pediatric population. *J Clin Endocrin Metabolism*, March 2015
- 47 Povisen J, Samonson MH. Fecal calprotectin in healthy children from 0 to 4 years, *ImmunoDiagnostics* no7, 2013
- 48 Immulite 2000 method sheets, Siemens
- 49 Alsultan A and Peloquin CA. Therapeutic drug monitoring in the treatment of tuberculosis: an update. *Drugs* 2014;74:839-54
- 50 Pugin J, Meisner M et al, Guide for the Clinical Use of PCT In Diagnosis and Monitoring of Sepsis April 2007
- 51 Kroonilise südamepuudulikkusega täiskasvanud patsiendi käsitus esmatasandil. *Eesti Ravijuhend. RJ-I/52.1-2021*
- 52 Roche cobas Valproehape – vaba valproehape rakendus. 2014-10 v5 Eesti keel.

Tähis	TÜLKHP-9.2.5
Viide	PÜL-09
Versioon	22